

Coupling Constants Evolution in a Universal Seesaw Mass Matrix Model

Yoshio Koide*

Department of Physics, University of Shizuoka

52-1 Yada, Shizuoka 422-8526, Japan

(February 1, 2008)

Abstract

Stimulated by a recent development of the universal seesaw mass matrix model, the evolutions of the gauge and Yukawa coupling constants are investigated under the gauge symmetries $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$. Especially, an investigation is made as to whether this evolution can constrain the necessary intermediate scales in these types of models and its viability.

12.15.Ff, 11.10.Hi, 12.60.-i

Typeset using REVTeX

*E-mail: koide@u-shizuoka-ken.ac.jp

I. INTRODUCTION

Recently, the so-called “universal seesaw mass matrix model” [1] has been revived [2,3] as a model which gives a unified description of masses and mixings of the quarks and leptons. The “seesaw mechanism” was first proposed [4] in order to answer the question of why neutrino masses are so invisibly small. Then, in order to understand that the observed quark and lepton masses are considerably smaller than the electroweak scale $\Lambda_L = \langle \phi_L^0 \rangle = 174 \text{ GeV}$, the mechanism was applied to the quarks [1]. However, the observation of the top quark of 1994 [5] aroused a doubt on the validity of the seesaw mechanism for the quarks because the observed fact $m_t \sim \Lambda_L$ means that $M_F^{-1}m_R$ is of the order of one in the seesaw expression $M_f \simeq m_L M_F^{-1} m_R$. On the contrary, it has recently been found [2,3] that the model can give an interpretation for the question of why only top quark acquires a mass of the order of Λ_L if we take an additional condition $\det M_F = 0$ for up-quark sector.

In the universal seesaw mass matrix model, the mass matrix for fermions (f, F) is given by

$$M = \begin{pmatrix} 0 & m_L \\ m_R & M_F \end{pmatrix} = m_0 \begin{pmatrix} 0 & Z_L \\ \kappa Z_R & \lambda Y_F \end{pmatrix}, \quad (1.1)$$

where f_i (fermion sector names $f = u, d, \nu, e$; family numbers $i = 1, 2, 3$) denote quarks and leptons, F_i denote hypothetical heavy fermions $F = U, D, N$ and E correspondingly to $f = u, d, \nu$ and e , and they belong to $f_L = (2, 1)$, $f_R = (1, 2)$, $F_L = (1, 1)$ and $F_R = (1, 1)$ of $\text{SU}(2)_L \times \text{SU}(2)_R$. The matrices Z_L , Z_R and Y_F are those of the order of one. The 3×3 matrices m_L ($\sim m_0 = \Lambda_L$) and m_R ($\sim \kappa m_0 = \Lambda_R$) are symmetry breaking mass terms of $\text{SU}(2)_L$ and $\text{SU}(2)_R$, respectively, and those have common structures independently of the fermion sector names f . Only M_F ($\sim \lambda m_0 = \Lambda_S$) has a structure dependent on the sector name f . For the case $\lambda \gg \kappa \gg 1$, the mass matrix (1.1) leads to the well-known seesaw expression

$$M_f \simeq m_L M_F^{-1} m_R. \quad (1.2)$$

In contrast to the case (1.2), for the case with the additional condition

$$\det M_F = 0 , \quad (1.3)$$

on the up-quark sector ($F = U$), one of the heavy fermions F_i (say, F_3) cannot acquire a mass of the order of $\Lambda_S \equiv \lambda m_0$, so that the seesaw mechanism does not work for the third fermion. Therefore, the mass generation at each energy scale is as follows: First, at the energy scale $\mu = \Lambda_S$, the heavy fermions F , except for U_3 , acquire the masses of the order of Λ_S . Second, at the energy scale $\mu = \Lambda_R$, the $SU(2)_R$ symmetry is broken, and the fermion u_{R3} generates a mass term of the order of Λ_R by pairing with U_{L3} . Finally, at $\mu = \Lambda_L$, the $SU(2)_L$ symmetry is broken, and the fermion u_{L3} generates a mass term of the order Λ_L by pairing with U_{R3} . The other fermions f acquire the well-known seesaw masses (1.2). The scenario is summarized in Table I. We regard the fermion pair (u_{L3}, U_{R3}) as the top-quark state. Thus, we can understand why only top quark t acquires the mass $m_t \sim O(m_L)$ [2,3].

On the other hand, for the neutrino mass generation, at present, we have the following two scenarios as summarized in Table II. One (Scenario A) is a trivial extension of the present model: we introduce a further large energy scale $\Lambda_{\nu S}$ in addition to Λ_S , and we assume that $M_F \sim \Lambda_S$ ($F = U, D, E$), while $M_N \sim \Lambda_{\nu S}$ ($\Lambda_{\nu S} \gg \Lambda_S$). Another scenario (Scenario B) [6] is one without introducing such an additional energy scale. The neutral heavy leptons are singlets of $SU(2)_L \times SU(2)_R$ and they do not have $U(1)$ -charge. Therefore, it is likely that they acquire Majorana masses M_M together with the Dirac masses $M_D \equiv M_N$ at $\mu = \Lambda_S$. For example, we assume $M_M = M_D$ [7]. Then, the neutrino mass matrix for the conventional light neutrinos is given by $M_\nu = m_L M_N^{-1} m_L^T$, so that the masses m_ν are given with the order of

$$m_\nu \sim \frac{\Lambda_L^2}{\Lambda_S} = \frac{1}{\kappa} \frac{\Lambda_L \Lambda_R}{\Lambda_S} . \quad (1.4)$$

In order to explain the smallness of m_ν , the model requires that the scale Λ_R must be extremely larger than Λ_L (for example, $\kappa \equiv \Lambda_R/\Lambda_L \sim 10^9$ [7]). This scenario seems to be very attractive from the theoretical point of view, because we can explain the mass hierarchy

of the quarks and leptons by the three energy scales Λ_L , Λ_R and Λ_S only. On the other hand, in the scenario A, there is no constraint on the value of κ (however, the value must be larger than ~ 10 because of no observation of the right-handed weak bosons W_R at present), so that the model allows a case with a lower value of Λ_R . Since we can expect abundant new physics effects for the case of $\kappa \sim 10$ [8], the case is also attractive from the phenomenological point of view.

One of the purposes of the present paper is to see whether a study of the evolutions of the gauge coupling constants of $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$ and of the Yukawa coupling constants in the universal seesaw mass matrix model can give any hint on the value of the intermediate energy scale Λ_R or not. For example, Shafi and Wetterich [9] and Rajpoot [10] have considered an $O(10)$ model and an $SO(10)$ model, respectively, with the symmetry breakings $SO(10) \rightarrow SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$ at $\mu = \Lambda_{GUT}$ and $SU(2)_R \rightarrow U(1)_R$ at $\mu = \Lambda_R$, and they have demonstrated that the model with $\Lambda_{GUT} \sim 10^{19}$ GeV and $\Lambda_R \sim 10^9$ GeV is consistent with the low energy phenomenology. The value $\Lambda_R \sim 10^9$ GeV is favorable to the scenario B for neutrino masses. However, in the present model, since there are many new fermions F above the intermediate energy scale Λ_S , their conclusion cannot be applied to the present seesaw mass matrix model straightforwardly.

On the other hand, a phenomenological study of the universal seesaw mass matrix model for the quark masses and the Cabibbo-Kobayashi-Maskawa (CKM) [11] matrix parameters has successfully been given by Fusaoka and the author [2]. In order to give explicit numerical predictions, they have used some working hypotheses that I will use here as well.

(i) The matrices Z_L and Z_R , which are universal for quarks and leptons, have the same structure:

$$Z_L = Z_R \equiv Z = \text{diag}(z_1, z_2, z_3) \quad , \quad (1.5)$$

with $z_1^2 + z_2^2 + z_3^2 = 1$, where, for convenience, we have taken a basis on which the matrix Z is diagonal.

(ii) The matrices Y_F , which have structures dependent on the fermion sector $f = u, d, \nu, e$,

take a simple form [(unit matrix)+(a rank one matrix)]:

$$Y_f = \mathbf{1} + 3b_f X \quad . \quad (1.6)$$

(iii) The rank one matrix X is given by a democratic form

$$X = \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} , \quad (1.7)$$

on the family-basis where the matrix Z is diagonal.

(iv) In order to fix the parameters z_i , we tentatively take $b_e = 0$ for the charged lepton sector, so that the parameters z_i are given by

$$\frac{z_1}{\sqrt{m_e}} = \frac{z_2}{\sqrt{m_\mu}} = \frac{z_3}{\sqrt{m_\tau}} = \frac{1}{\sqrt{m_e + m_\mu + m_\tau}} \quad . \quad (1.8)$$

By taking $b_u = -1/3$ (then $\det M_U = 0$), they have obtained the following top-quark mass enhancement without the suppression factor κ/λ

$$m_t \simeq \frac{1}{\sqrt{3}} m_0 , \quad (1.9)$$

together with the successful relation $m_u/m_c \simeq 3m_e/4m_\mu$. Furthermore, by taking $b_d = -e^{i\beta_d}$ ($\beta_d = 18^\circ$), they have succeeded in giving the reasonable values of the CKM matrix parameters together with the reasonable values of the quark mass ratios (not only m_i^u/m_j^u , m_i^d/m_j^d , but also m_i^u/m_j^d) with keeping the value of the parameter $(m_0\kappa/\lambda)_f$ in $(m_0\kappa/\lambda)_u = (m_0\kappa/\lambda)_d$. However, in order to fit the quark mass values (not the ratios) to the observed quark mass values at $\mu = m_Z$, they have taken the parameter $(m_0\kappa/\lambda)_f$ as

$$R(m_Z) \equiv \left(\frac{(m_0\kappa/\lambda)_u}{(m_0\kappa/\lambda)_e} \right)_{\mu=m_Z} \simeq 3 \quad . \quad (1.10)$$

It seems to be natural to consider that all Yukawa coupling constants become equal between quarks and leptons at a large energy scale Λ_{YU} . Therefore, another one of the purposes of the present paper is to see whether such a factor 3 can be understood by the difference of the

evolutions of the Yukawa coupling constants between quarks and leptons from the energy scale $\mu = \Lambda_{YU}$ to $\mu = m_Z$.

In Sec. II and Sec. III, we investigate evolution of the gauge and Yukawa coupling constants, respectively, under the gauge symmetries $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$ at one-loop. We will conclude that it is possible to find the energy scale Λ_{YU} at which $R(\mu)$ takes $R = 1$ only for a model with a value $\kappa < 10^2$. Although in Sec. II and Sec. III we consider the case that the symmetries $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$ are unbroken for the region $\mu > \Lambda_S$, in Sec. IV, we investigate a case that the symmetries $SU(3)_c \times U(1)_Y$ are embedded into the Pati-Salam symmetry [12] $SU(4)_{PS}$ at $\mu > \Lambda_S$, so that we consider the case of $SU(4)_{PS} \times SU(2)_L \times SU(2)_R$ in the region $\Lambda_S < \mu \leq \Lambda_{GUT}$. We will find that the model predicts $\Lambda_R \simeq 5 \times 10^{12}$ GeV, $\Lambda_S \simeq 3 \times 10^{14}$ GeV, and $\Lambda_{GUT} \simeq 6 \times 10^{17}$ GeV. Finally, Sec. V is devoted to the conclusions and remarks. We will find that there is no model which satisfies $\Lambda_{YU} = \Lambda_{GUT}$.

II. EVOLUTION OF THE GAUGE COUPLING CONSTANTS

The gauge symmetries $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$ are broken into the gauge symmetries $SU(3)_c \times SU(2)_L \times U(1)_{Y'}$ at $\mu = \Lambda_R$. The electric charge operator Q

$$Q = I_3^L + I_3^R + \frac{1}{2}Y, \quad (2.1)$$

at $\mu > \Lambda_R$ is changed into

$$Q = I_3^L + \frac{1}{2}Y', \quad (2.2)$$

in the region $\Lambda_L < \mu \leq \Lambda_R$. Hereafter, we call the regions $\Lambda_L < \mu \leq \Lambda_R$, $\Lambda_R < \mu \leq \Lambda_S$ and $\Lambda_S < \mu \leq \Lambda_X$ ($\Lambda_X \equiv \Lambda_{GUT}$ or $\Lambda_X \equiv \Lambda_{YU}$) Regions I, II and III, respectively.

The evolutions of the gauge coupling constants g_i at one-loop are given by the equations

$$\frac{d}{dt}\alpha_i(\mu) = -\frac{1}{2\pi}b_i\alpha_i^2(\mu), \quad (2.3)$$

where $\alpha_i \equiv g_i^2/4\pi$, $t = \ln \mu$ and the coefficients b_i are given in Table IV. (Note that the heavy fermions F_L and F_R except for U_{L3} and U_{R3} are decoupled for $\mu \leq \Lambda_S$ and the fermions u_{R3} and U_{L3} are decoupled for $\mu \leq \Lambda_R$.) The boundary conditions at $\mu = \Lambda_L$ and $\mu = \Lambda_R$ are as follows:

$$\alpha_{em}^{-1}(\Lambda_L) = \alpha_L^{-1}(\Lambda_L) + \frac{5}{3}\alpha_1'^{-1}(\Lambda_L) , \quad (2.4)$$

and

$$\frac{5}{3}\alpha_1'^{-1}(\Lambda_R) = \alpha_R^{-1}(\Lambda_R) + \frac{2}{3}\alpha_1^{-1}(\Lambda_R) , \quad (2.5)$$

respectively, correspondingly to (2.2) and (2.1), where the normalizations of the $U(1)_{Y'}$ and $U(1)_Y$ gauge coupling constants have been taken as they satisfy $\alpha_1' = \alpha_L = \alpha_3$ in the $SU(5)$ grand-unification limit and $\alpha_1 = \alpha_L = \alpha_R = \alpha_3$ in the $SO(10)$ grand-unification limit, respectively. For convenience, we use the initial values at $\mu = m_Z$ instead of those at $\mu = \Lambda_L$ in the region I: $\alpha_1'(m_Z) = 0.01683$, $\alpha_L(m_Z) = 0.03349$ and $\alpha_3(m_Z) = 0.118$. The values of α_1' and α_L have been derived from [13] $\alpha_{em}(m_Z) = (128.89 \pm 0.09)^{-1}$ and $\sin^2 \theta_W = 0.23165 \pm 0.00024$. The value of α_3 has been quoted from Ref. [14]. We illustrate a typical case with $\Lambda_R = 10^5$ GeV and $\alpha_R(\Lambda_R) = 1/4\pi$ in Fig. 1.

In the numerical study, we have taken the value of the parameter $\Lambda_R/\Lambda_S \equiv \kappa/\lambda$ as

$$\kappa/\lambda = \Lambda_R/\Lambda_S = 0.02 , \quad (2.6)$$

which has been obtained from the observed value of the ratio m_c/m_t in Ref. [2]. Although the value (2.6) has been obtained on the model with the specific matrix forms (1.5) - (1.7), the order of the value (2.6) will be valid for any other seesaw model with $\det M_U = 0$ because in such a model the value of κ/λ is given by the order of m_b/m_t .

As seen in Fig. 1, the $U(1)$ coupling constant $\alpha_1(\mu)$ becomes rapidly strong in the region III ($\mu > \Lambda_S$) because the heavy fermions F become massless in the region III. We consider that the unification energy scale Λ_{YU} of the Yukawa coupling constants must be lower than an energy scale Λ_1^∞ at which $\alpha_1(\mu)$ becomes infinity. This condition will impose a strong

restriction on the possible Λ_{YU} -search as we discuss in the next section. Of course, in the grand unification scenario, the $U(1)$ symmetry will be embedded into a grand unification symmetry G before the $U(1)$ coupling constant bursts. Such a case will be discussed in Sec. IV.

III. EVOLUTION OF $y_L y_R / y_S$

The 3×3 matrices m_L , m_R and M_F are given in terms of the vacuums expectation values $v_L = \sqrt{2}\langle\phi_L^0\rangle$, $v_R = \sqrt{2}\langle\phi_R^0\rangle$ and $v_S = \langle\Phi\rangle$, and the matrices Z and Y_F defined by (1.5) - (1.7) as follows:

$$m_L^f = \frac{1}{\sqrt{2}} y_L^f v_L Z, \quad m_R^f = \frac{1}{\sqrt{2}} y_R^f v_R Z, \quad M_F = y_S^f v_S Y_F. \quad (3.1)$$

The evolution of the Yukawa coupling constants are given by

$$\frac{d}{dt} \ln(y_L^f Z) = \frac{1}{16\pi^2} (T_L^f - G_L^f + H_L^f), \quad (3.2)$$

$$\frac{d}{dt} \ln(y_R^f Z) = \frac{1}{16\pi^2} (T_R^f - G_R^f + H_R^f), \quad (3.3)$$

$$\frac{d}{dt} \ln(y_S^f Y_F) = \frac{1}{16\pi^2} (T_S^f - G_S^f + H_S^f), \quad (3.4)$$

where T^f , G^f and H^f denote contributions from fermion-loop corrections, vertex corrections due to the gauge bosons and vertex corrections due to the Higgs boson, respectively.

What is of great interest to us is to see whether the evolutions can explain the value $R(m_Z) \simeq 3$ or not, i.e., our interest exists not in the hierarchy among m_e , m_μ and m_τ , but in the hierarchy among up-quark, down-quark, charged lepton and neutrino sectors. Therefore, we neglect the scale-dependency of the matrix Z , because we can regard the value of z_3 as $z_3 \simeq 1$ from Eq. (1.8). We also neglect the scale-dependency of the matrix Y_F because the matrices Y_F are expressed as

$$Y_F = \text{diag}(1, 1, 1 + 3b_f), \quad (3.5)$$

on the basis on which the matrix Y_F is diagonal, and we find that the forms $Y_E = \text{diag}(1, 1, 1)$ and $Y_U = \text{diag}(1, 1, 0)$ are scale-invariant and $Y_D = \text{diag}(1, 1, 1 - 3e^{i\beta_d})$ is almost scale-invariant. For convenience, we approximately still use the evolution equation of $y_S Y_F$ at $\mu \leq \Lambda_S$ and that of $y_R Z$ at $\mu \leq \Lambda_R$. Then, the ratio $R(\mu)$ defined by Eq. (1.10) can be expressed in terms of the Yukawa coupling constants y_L , y_R and y_S as follow:

$$R(\mu) \equiv \frac{y_L^u(\mu)y_R^u(\mu)/y_S^u(\mu)}{y_L^e(\mu)y_R^e(\mu)/y_S^e(\mu)} . \quad (3.6)$$

The evolution of the ratio $R(\mu)$ is approximately given by

$$\frac{d}{dt} \ln R(\mu) \simeq -\frac{1}{16\pi^2} (G - H) , \quad (3.7)$$

where

$$\begin{aligned} G &= (G_L^u + G_R^u - G_S^u) - (G_L^e + G_R^e - G_S^e) , \\ H &= (H_L^u + H_R^u - H_S^u) - (H_L^e + H_R^e - H_S^e) . \end{aligned} \quad (3.8)$$

The G - and H -terms are given in Table V. Since in the present model, $|y_L^u|^2 \simeq |y_L^d|^2$, $|y_L^e|^2 \simeq |y_L^\nu|^2$, and so on, differently from other models where $|y_L^d/y_L^u| \simeq m_b/m_t$ and $|y_L^\nu/y_L^e| \simeq m_\nu/m_\tau$, we can neglect the H_L - and H_R -terms in Eq. (3.8). When we also neglect the H_S -terms, the ratio $R(\mu)$ is approximately evaluated as follows:

$$\frac{R(\mu)}{R(m_Z)} = \left(1 + \frac{b_3^I}{2\pi} \alpha_3(m_Z) \ln \frac{\mu}{m_Z}\right)^{-4/b_3^I} \left(1 + \frac{b_1^I}{2\pi} \alpha_1'(m_Z) \ln \frac{\mu}{m_Z}\right)^{7/10b_1^I} , \quad (3.9)$$

$$\frac{R(\mu)}{R(\Lambda_R)} = \left(1 + \frac{b_3^{II}}{2\pi} \alpha_3(\Lambda_R) \ln \frac{\mu}{\Lambda_R}\right)^{-4/b_3^{II}} \left(1 + \frac{b_1^{II}}{2\pi} \alpha_1(\Lambda_R) \ln \frac{\mu}{\Lambda_R}\right)^{1/b_1^{II}} , \quad (3.10)$$

$$\frac{R(\mu)}{R(\Lambda_S)} = \left(1 + \frac{b_3^{III}}{2\pi} \alpha_3(\Lambda_S) \ln \frac{\mu}{\Lambda_S}\right)^{-4/b_3^{III}} \left(1 + \frac{b_1^{III}}{2\pi} \alpha_1(\Lambda_R) \ln \frac{\mu}{\Lambda_S}\right)^{1/b_1^{III}} , \quad (3.11)$$

for the regions I, II and III, respectively. By using (3.9) - (3.11), we can obtain the energy scale $\mu = \Lambda_{YU}$ at which the ratio $R(\mu)$ takes $R(\Lambda_{YU}) = 1$.

In Fig. 2, we illustrate the behavior of Λ_{YU} for a given value of Λ_R . For reference, we also illustrate the behavior of Λ_1^∞ , at which $\alpha_1^{-1}(\mu)$ takes $\alpha_1^{-1}(\Lambda_1^\infty) = 0$. The value of Λ_{YU} must be lower than the value of Λ_1^∞ . Therefore, as seen in Fig. 2, if we adhere to the constraint $\alpha_R(\Lambda_R) = \alpha_L(\Lambda_R)$, we must abandon a model with a higher κ value ($\kappa > 10^2$). Only a model with $\kappa \sim 10$ is acceptable. However, if we admit a strong coupling of the right-handed weak bosons at $\mu = \Lambda_R$, for example, $\alpha_R(\Lambda_R) \geq 1/4\pi$, a model with a higher κ value also becomes acceptable.

Of course, from a similar study, we can find that the evolution of $R_{u/d}(\mu) \equiv (y_L^u y_R^u / y_S^u) / (y_L^d y_R^d / y_S^d)$ still keeps $R_{u/d}(m_Z) \simeq 1$. Therefore, the parametrization $(m_0 \kappa / \lambda)_u = (m_0 \kappa / \lambda)_d$ in Ref. [2] is justified.

IV. EVOLUTION OF THE PATI-SALAM COLOR

In order to avoid the burst of the U(1) gauge coupling constant, we consider that the $U(1)_Y \times SU(3)_c$ symmetries are embedded into the Pati-Salam $SU(4)$ symmetry [12] above $\mu = \Lambda_S$. In other words, the $SU(4)_{PS}$ gauge symmetry is broken into $SU(3)_c \times U(1)_Y$ at $\mu = \Lambda_S$. Indeed, the structures of the heavy fermion mass matrices M_F are flavor-dependent. The fermions f and F belong to $f_L = (2, 1, 4)$, $f_R = (1, 2, 4)$, $F_L = (1, 1, 4)$ and $F_R = (1, 1, 4)$ of $SU(2)_\times SU(2)_R \times SU(4)_{PS}$ at $\mu \geq \Lambda_S$.

In the region III ($\Lambda_S < \mu \leq \Lambda_X$), $\alpha_L(\mu)$ and $\alpha_R(\mu)$ are evolved with the coefficients b_L^{III} and b_R^{III} given in Table IV, but $\alpha_1(\mu)$ and $\alpha_3(\mu)$ are replaced with $\alpha_4(\mu)$ which is evolved with the coefficient

$$b_4^{III} = 20/3, \quad (4.1)$$

where the boundary condition at $\mu = \Lambda_S$ is

$$\alpha_1(\Lambda_S) = \alpha_3(\Lambda_S) = \alpha_4(\Lambda_S). \quad (4.2)$$

Since $\alpha_1(\Lambda_S)$ and $\alpha_3(\Lambda_S)$ are given by

$$\alpha_1^{-1}(\Lambda_S) = \frac{5}{2} \left[\alpha_1^{-1}(\Lambda_L) + \frac{b_1^I}{2\pi} \ln \frac{\Lambda_R}{\Lambda_L} \right] - \frac{3}{2} \left[\alpha_L^{-1}(\Lambda_L) + \frac{b_L^I}{2\pi} \ln \frac{\Lambda_R}{\Lambda_L} \right] + \frac{b_1^{II}}{2\pi} \ln \frac{\Lambda_S}{\Lambda_R} , \quad (4.3)$$

$$\alpha_3^{-1}(\Lambda_S) = \alpha_3^{-1}(\Lambda_L) + \frac{b_3^I}{2\pi} \ln \frac{\Lambda_R}{\Lambda_L} + \frac{b_3^{II}}{2\pi} \ln \frac{\Lambda_S}{\Lambda_R} , \quad (4.4)$$

respectively, the values of Λ_R and Λ_S are fixed at

$$\Lambda_R = 5.46 \times 10^{12} \text{ GeV} , \quad \Lambda_S = 2.73 \times 10^{14} \text{ GeV} , \quad (4.5)$$

under the conditions (2.6) and (4.2). The unification scale Λ_{GUT} is also fixed at

$$\Lambda_{GUT} = 5.84 \times 10^{17} \text{ GeV} , \quad (4.6)$$

by the condition

$$\alpha_L(\Lambda_{GUT}) = \alpha_R(\Lambda_{GUT}) = \alpha_4(\Lambda_{GUT}) , \quad (4.7)$$

for example, for the embedding into $\text{SO}(10)$ [8,9]. In Fig. 3, we illustrate the behaviors of $\alpha_i^{-1}(\mu)$. Roughly speaking, the value $\Lambda_R \sim 10^{12}$ is favorable to the scenario B for the neutrino mass generation.

On the other hand, the evolution of $R(\mu)$ defined by (3.6) is almost constant at $\mu \geq \Lambda_S$, i.e., $R(\Lambda_S) \simeq R(\Lambda_{YU})$, because there is no difference between quarks and leptons in the region III ($\Lambda_S < \mu \leq \Lambda_{YU}$). Therefore, we obtain

$$R(m_Z)/R(\Lambda_{YU}) \simeq R(m_Z)/R(\Lambda_S) = 2.3 , \quad (4.8)$$

and we fail to obtain our desirable relation $R(m_Z)/R(\Lambda_{YU}) \simeq 3$. If we adhere the unification of the gauge symmetries $\text{SU}(3)_c \times \text{SU}(2)_L \times \text{SU}(2)_R \times \text{U}(1)_Y$ into a Pati-Salam type unification G , we must abandon the idea that the discrepancy $R(m_Z) \simeq 3$ between quarks and leptons in the model given in Ref. [2] comes from difference of evolutions between quarks and leptons, or we must consider that the magnitudes of the Yukawa coupling constants are different between quarks and leptons from the beginning at $\mu = \Lambda_{GUT}$.

V. CONCLUDING REMARKS

In conclusion, we have investigated the evolution of the universal seesaw mass matrix model under the gauge symmetries $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_Y$. The symmetries can be embedded into a unification symmetry of the Pati-Salam type at $\Lambda_{GUT} = 5.84 \times 10^{17}$ GeV. Then, the value $\Lambda_R = 5.46 \times 10^{12}$ GeV is favorable to the scenario B for the neutrino mass generation. However, we cannot explain the discrepancy $R(m_Z) \simeq 3$ between quarks and leptons by the evolution of $R(\mu)$ starting from $R(\Lambda_{GUT}) = 1$.

On the other hand, if we abandon the grand unification scenario, the model has a possibility that the value $R(m_Z) \simeq 3$ can be understood by the evolution of the Yukawa coupling constants. As seen in Fig. 2, we require $\alpha_L(\Lambda_R) = \alpha_R(\Lambda_R)$, the value Λ_R for the case which gives $R(m_Z) \simeq 3$ must be $\Lambda_R \leq 10^4$ GeV. If we accept a model with a strong $SU(2)_R$ force at $\mu = \Lambda_R$, for example, $\alpha_R(\Lambda_R) = 1/4\pi$, the region $\Lambda_R \leq 10^{18}$ GeV also becomes allowed. We consider that the model with $\kappa \sim 10$ is likely. Although this case rules out the scenario B for neutrinos, phenomenologically we can expect an abundance of new physics effects [8], t' production, FCNC effects, and so on, in the near future colliders.

Our numerical results have been obtained by the constraint $\Lambda_R/\Lambda_S = 0.02$ which has come from the observed ratio of m_c/m_t under the special model with (1.5) - (1.7). Since the ratio Λ_R/Λ_S is fixed by the ratio m_c/m_t (or m_b/m_t) as far as a universal seesaw model with $\det M_U = 0$ are concerned, the value of the ratio Λ_R/Λ_S is, in general, of the order of 10^{-2} . Therefore, our conclusions will be unchanged as far as the orders are concerned.

In the present paper, we have not discussed a SUSY version of the present model, although the case is attractive from the point of view of the grand unification. In such a SUSY version, since the coefficient $8\alpha_3$ in G -terms in (3.8) [also in Table V] is changed for $(16/3)\alpha_3$, the case push the energy scale Λ_{YU} to an unlikely ultra-high energy scale ($> 10^{23}$ GeV). If we want to adopt a SUSY version of the present model, we must abandon the idea of the unification of the Yukawa coupling constants.

ACKNOWLEDGMENTS

The author would like to thank M. Tanimoto, T. Matsuoka and N. Okamura for their helpful comments, especially, on the evolution equations. This work was supported by the Grand-in-Aid for Scientific Research, Ministry of Education, Science and Culture, Japan (No. 08640386).

REFERENCES

- [1] Z. G. Berezhiani, Phys. Lett. **129B**, 99 (1983); Phys. Lett. **150B**, 177 (1985); D. Chang and R. N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); A. Davidson and K. C. Wali, Phys. Rev. Lett. **59**, 393 (1987); S. Rajpoot, Mod. Phys. Lett. **A2**, 307 (1987); Phys. Lett. **191B**, 122 (1987); Phys. Rev. **D36**, 1479 (1987); K. B. Babu and R. N. Mohapatra, Phys. Rev. Lett. **62**, 1079 (1989); Phys. Rev. **D41**, 1286 (1990); S. Ranfone, Phys. Rev. **D42**, 3819 (1990); A. Davidson, S. Ranfone and K. C. Wali, Phys. Rev. **D41**, 208 (1990); I. Sogami and T. Shinohara, Prog. Theor. Phys. **66**, 1031 (1991); Phys. Rev. **D47**, 2905 (1993); Z. G. Berezhiani and R. Rattazzi, Phys. Lett. **B279**, 124 (1992); P. Cho, Phys. Rev. **D48**, 5331 (1994); A. Davidson, L. Michel, M. L. Sage and K. C. Wali, Phys. Rev. **D49**, 1378 (1994); W. A. Ponce, A. Zepeda and R. G. Lozano, Phys. Rev. **D49**, 4954 (1994).
- [2] Y. Koide and H. Fusaoka, Z. Phys. **C71**, 459 (1996); Prog. Theor. Phys. **97**, 459 (1997).
- [3] T. Morozumi, T. Satou, M. N. Rebelo and M. Tanimoto, Phys. Lett. **B410**, 233 (1997).
- [4] M. Gell-Mann, P. Rammond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, 1979); T. Yanagida, in *Proc. Workshop of the Unified Theory and Baryon Number in the Universe*, edited by A. Sawada and A. Sugamoto (KEK, 1979); R. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [5] CDF collaboration, F. Abe *et al.*, Phys. Rev. Lett. **73**, 225 (1994).
- [6] Z. G. Berezhiani, in Ref.[1]; A. Davidson and K. C. Wali, in Ref.[1]; S. Rajpoot, in Ref.[1]; A. Davidson, S. Ranfone and K. C. Wali, in Ref.[1]; W. A. Ponce, A. Zepeda and R. G. Lozano, in Ref.[1].
- [7] Y. Koide, hep-ph/9707505 (1997), to be published in Phys. Rev. **D**, No.9 (1998).
- [8] Y. Koide, Phys. Rev. **D56**, 2656 (1997).

- [9] Q. Shafi and C. Wetterich, Phys. Lett. **85B**, 52 (1979).
- [10] S. Rjpoot, Phys. Rev. **D22**, 2244 (1980).
- [11] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1996); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [12] J. C. Pati and A. Salam, Phys. Rev. Lett. **31**, 661 (1973); **36**, 1229 (1976); Phys. Lett. **58B**, 333 (1975); Phys. Rev. **D8**, 1240 (1973); **D10**, 275 (1974).
- [13] W. Hollik, University of Karlsruhe Report No. KA-TP-19-1996, 1996 (unpublished);
See also Z. Hioki, Acta Phys. Pol. **B27**, 1569 (1996).
- [14] I. Hinchliffe, Particle data group, Phys. Rev. **D54**, 77 (1996).

TABLES

TABLE I. Fermion mass generation scenario

Energy scale	d - & e -sectors	u -sector ($i \neq 3$)
At $\mu = \Lambda_S \sim \lambda m_0$	$m(F_L, F_R) \sim \Lambda_S$	$m(U_{Li}, U_{Ri}) \sim \Lambda_S$
At $\mu = \Lambda_R \sim \kappa m_0$		$m(u_{R3}, U_{L3}) \sim \Lambda_R$
At $\mu = \Lambda_L \sim m_0$		$m(u_{L3}, U_{R3}) \sim \Lambda_L$
	$m(f_L, f_R) \sim \frac{\Lambda_L \Lambda_R}{\Lambda_S}$	$m(u_{Li}, u_{Ri}) \sim \frac{\Lambda_L \Lambda_R}{\Lambda_S}$

TABLE II. Neutrino mass generation scenarios: $N_{\pm} = (N_L \pm N_R^c)/\sqrt{2}$

Energy scale	Scenario A	Scenario B
At $\mu = \Lambda_{\nu S}$	$m(N_L, N_R) \sim \Lambda_{\nu S}$	
At $\mu = \Lambda_S$		$m(N_+, N_+^c) \sim \Lambda_S$
At $\mu = \Lambda_R$		$m(\nu_R, N_-) \sim \Lambda_R$
At $\mu = \Lambda_L$	$m(\nu_L, \nu_R) \sim \frac{\Lambda_L \Lambda_R}{\Lambda_{\nu S}}$	$m(\nu_L, \nu_L^c) \sim \frac{\Lambda_L^2}{\Lambda_S}$

TABLE III. Quantum numbers of the fermions f and F and Higgs scalars ϕ_L , ϕ_R and Φ .

	I_3^L	I_3^R	Y		I_3^L	I_3^R	Y
u_L	$+\frac{1}{2}$	0	$\frac{1}{3}$	u_R	0	$+\frac{1}{2}$	$\frac{1}{3}$
d_L	$-\frac{1}{2}$	0	$\frac{1}{3}$	d_R	0	$-\frac{1}{2}$	$\frac{1}{3}$
ν_L	$+\frac{1}{2}$	0	-1	ν_R	0	$+\frac{1}{2}$	-1
e_L	$-\frac{1}{2}$	0	-1	e_R	0	$-\frac{1}{2}$	-1
U_L	0	0	$\frac{4}{3}$	U_R	0	0	$\frac{4}{3}$
D_L	0	0	$-\frac{2}{3}$	D_R	0	0	$-\frac{2}{3}$
N_L	0	0	0	N_R	0	0	0
E_L	0	0	-2	E_R	0	0	-2
ϕ_L^+	$+\frac{1}{2}$	0	1	ϕ_R^+	0	$+\frac{1}{2}$	1
ϕ_L^0	$-\frac{1}{2}$	0	1	ϕ_R^0	0	$-\frac{1}{2}$	1
Φ	0	0	0				

TABLE IV. Coefficients in the evolution equations of gauge coupling constants.

	$\Lambda_L < \mu \leq \Lambda_R$	$\Lambda_R < \mu \leq \Lambda_S$	$\Lambda_S < \mu \leq \Lambda_X$
$SU(3)_c$	$b_3^I = 7$	$b_3^{II} = 19/3$	$b_3^{III} = 3$
$SU(2)_L$	$b_L^I = 19/6$	$b_L^{II} = 19/6$	$b_L^{III} = 19/6$
$SU(2)_R$		$b_R^{II} = 19/6$	$b_R^{III} = 19/6$
$U(1)_Y$	$b_1^I = -41/10$	$b_1^{II} = -43/6$	$b_1^{III} = -41/2$

TABLE V. G - and H -terms in the evolution equation (3.8). H_L^e , H_R^e and H_S^e are given by the replacements $u \rightarrow e$ and $d \rightarrow \nu$ in H_L^u , H_R^u and H_S^u , respectively.

	$I \ (\Lambda_L < \mu \leq \Lambda_R)$	$II \ (\Lambda_R < \mu \leq \Lambda_S)$	$III \ (\Lambda_S < \mu \leq \Lambda_{YU})$
$G =$	$4\pi \left(8\alpha_3 - \frac{7}{5}\alpha'_1\right)$	$4\pi (8\alpha_3 - 2\alpha_1)$	
$H_L^u =$	$\frac{3}{2} \left(y_L^u ^2 - y_L^d ^2\right)$		
$H_R^u =$	0	$\frac{3}{2} \left(y_R^u ^2 - y_R^d ^2\right)$	
$H_S^u =$	0		$3 y_S^u ^2$

FIGURES

FIG. 1. Behaviors of $\alpha_3^{-1}(\mu)$ (dotted chain line), $\alpha_L^{-1}(\mu)$ (dotted line), $\alpha_R^{-1}(\mu)$ (broken line) and $\alpha_1^{-1}(\mu)$ [$\alpha_1'^{-1}(\mu)$] (solid line) in the case with the input values $\Lambda_R = 10^5$ GeV and $\alpha_R(\Lambda_R) = 1/4\pi$.

FIG. 2. Behavior of Λ_{YU} versus Λ_R . The bold and thin solid lines denote the cases of the input values $\alpha_R(\Lambda_R) = \alpha_L(\Lambda_R)$ and $\alpha_R(\Lambda_R) = 1/4\pi$, respectively. For reference, the behaviors of Λ_1^∞ for the cases of the input values $\alpha_R(\Lambda_R) = \alpha_L(\Lambda_R)$ (bold broken line) and $\alpha_R(\Lambda_R) = 1/4\pi$ (thin broken line) are illustrated. The physical value of Λ_{YU} must be $\Lambda_{YU} < \Lambda_1^\infty$.

FIG. 3. Behaviors of $\alpha_3^{-1}(\mu)$ [$\alpha_4^{-1}(\mu)$ for $\Lambda_S < \mu \leq \Lambda_{GUT}$] (dotted chain line), $\alpha_L^{-1}(\mu)$ [= $\alpha_R^{-1}(\mu)$ for $\mu > \Lambda_R$] (dotted line) and $\alpha_1^{-1}(\mu)$ [$\alpha_1'^{-1}(\mu)$ for $\Lambda < \mu \leq \Lambda_R$] (solid line) in the case of the Pati-Salam type unification, where $\Lambda_R = 5.46 \times 10^{12}$ GeV, $\Lambda_S = 2.37 \times 10^{14}$ GeV and $\Lambda_{GUT} = 5.84 \times 10^{17}$ GeV.